

Clinical Study of Single Stage Dorsolateral Onlay Buccal Mucosal Graft Urethroplasty in Long Segment Anterior Urethral Stricture

Mondal PP¹, Das SK^{2*}, Mahapatra RS³ and Bera MK¹

¹Department of Urology, Kolkata Medical College, Kolkata, India

²Department of Urology, Burdwan Medical College, Burdwan, India

³Department of Urology, Bankura Medical College, Bankura, West Bengal, India

*Corresponding author:

Dr. Susanta Kumar Das,
Department of Urology, Burdwan Medical College,
Burdwan, West Bengal, India,
E-mail: susanta_kr_das@rediffmail.com

Received: 12 Jan 2022

Accepted: 21 Jan 2022

Published: 28 Jan 2022

J Short Name: COS

Copyright:

©2022 Das Sk. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Citation:

Das Sk, Clinical Study of Single Stage Dorsolateral Onlay Buccal Mucosal Graft Urethroplasty in Long Segment Anterior Urethral Stricture. Clin Surg. 2022; 7(1): 1-4

Keywords:

Stricture; Dorsolateral onlay urethroplasty; Buccal mucosal graft

1. Abstract

1.1. Objectives: To evaluate the results of dorsolateral onlay buccal mucosal graft (BMG) urethroplasty in various types of long segment anterior urethral stricture.

1.2. Materials and Methods: This is a prospective study, conducted in the Department of Urology, Medical College, Kolkata and Burdwan Medical College, Burdwan from April 2017 to March 2021. Forty patients with long segment (more than 1.5 cm) anterior urethral stricture who underwent substitution urethroplasty using dorsolateral onlay BMG urethroplasty were evaluated with detailed history, physical examination, ultrasonography of KUB with PVRU, uroflowmetry, retrograde urethrography and voiding cystourethrography, sono-urethrograms and cystoscopy. After treatment with dorsolateral onlay BMG urethroplasty, they were followed-up with uroflowmetry at 3 monthly interval for one year and RGU and/or urethroscopy if indicated. Success was defined as normal voiding pattern without any intervention post-operatively.

1.3. Results: With nineteen of the forty cases (47.5%), lichen sclerosis formed the single most common etiology. Panurethral strictures were more common. Out of total 40 cases, 33 (82.5%) had successful outcome and 7 (17.5%) were failure.

1.4. Conclusion: We conclude that it is a simple, technically feasible, easily adoptable and finally successful procedure.

2. Introduction

Urethral stricture disease with varied etiologies is common among men [1]. Management of anterior urethral stricture poses a continuing urological challenge.

The treatment of urethral stricture varies according to etiology, location, length, depth and density of the stricture (spongiofibrosis), multiplicity, proximity to sphincteric mechanism and the presence of local adverse factors.

There are different treatment options for short urethral stricture ranging from simple dilatation, directly visualized optical internal urethrotomy, resection and end to end anastomosis.

Long segment anterior urethral strictures that are not amenable to end-end anastomosis require substitution urethroplasty or may be managed by two-stage urethroplasty with or without use of free graft. Although augmented anastomotic techniques are currently suggested for these kind of strictures, the material for reconstruction (flap or graft) and location of the on the urethral surface (ventral or dorsal) has become a continuous issue [2-4].

Long segment anterior urethral stricture that require substitution urethroplasty uses penile skin flap, split and full thickness skin graft that can be derived from scrotum, penis, extragenital sites, tunica vaginalis, bladder mucosa, colonic mucosa, buccal mucosa or lingual mucosa.

Buccal mucosa graft becomes an ideal urethral substitute because contains no hair, flexible, easy to harvest, resistance to infection, compatibility with a wet environment, thick epithelium with a thin but well-vascularized lamina propria, allowing easy inosculation and survival.

For the management of anterior urethral stricture, dorsal, ventral or lateral onlay urethroplasty may be employed as single-stage procedure [5]. Out of these, dorsal onlay urethroplasty currently

most favoured [6]. We performed dorsolateral onlay urethroplasty by unilateral urethral mobilization [7, 8]. In dorsolateral approach instead of circumferential mobilization of urethra, mobilization of urethra is done in one side, i.e., from ventral midline to beyond dorsal midline. It preserves neuro-vascular integrity of one side of urethra thus reduce ischaemia with resultant chordee. Entire procedure is carried out by a single perineal incision and no incision is made on the penis except meatotomy when required.

Outcome of one stage transperineal dorsolateral onlay BMG urethroplasty was assessed in this study.

3. Materials and Methods

This prospective study was conducted in the departments of Urology, Medical College Hospital, Kolkata and Burdwan Medical College, Burdwan from April 2017 to March 2020 having a total case of 40 dorsolateral buccal mucosal graft urethroplasty.

Patients between the age group of 25 to 65 with anterior urethral stricture more than 1.5 cm without any previous history of urethral dilatation, Direct Visual Internal Urethrotomy (DVIU) or prior urethroplasty were included as study population.

Patients with severe co-morbidity, small stricture <1.5 cm, posterior urethral strictures, complete urethral blockade or stricture not allowing 6 Fr infant feeding tube were excluded from the study.

Preoperative evaluation included detailed history documenting international prostate symptom score (IPSS), prior operative intervention, history of extra-marital sexual exposure, recurrent urinary tract infection, urethral discharge, trauma and history of any acute urinary tract obstruction leading to supra-pubic cystostomy. In physical examination, we looked for circumcised status, any surgical scar and health of preputial skin and oral mucosa. Urinalysis, ultrasonography of KUB region with post-voided residual, retrograde urethrography, voiding cystourethrography, sonourethrogram, uroflowmetry, urethroscopy and other routine investigations were done before

Surgeries were done under general anaesthesia with nasotracheal intubation. Urethroplasty operations were performed in perineal approach, with the patients positioned in the exaggerated dorsal lithotomy position. All pressure points were appropriately padded with cotton. The penis was everted through the midline perineal incision. The bulbocavernous muscle was exposed and divided. The corpus spongiosum of the anterior urethra was exposed. Then the bulbar urethra was dissected to separate from corporacavernosa on left side, so as to leave the right half attached and preservation of its lateral blood supply. By invaginating the penis, the penile urethra was similarly dissected up to the coronal sulcus [9]. Then the strictured segment of the urethra was identified and opened dorsolaterally.

Buccal mucosal harvesting was done from one or both cheeks with or without lower lip, depending on the length of the stricture (Figure 1). The graft was then defatted, multiple opening done and tai-

lored to its proper size. The dissected urethra was rotated towards right side to expose the dorsolateral surface of the strictured segment of urethra and opened vertically extending the incision for about 1 cm both proximally and distally into the normal urethral lumen. The proximal and distal urethral lumen were calibrated. One edge of the graft was sutured to the margin fixed to the corpora with 4-0 polyglactin interrupted sutures. A 16 Fr silicone catheter was introduced. The other margin of the graft was sutured to the lateral margin of the urethra and the tunica of corporacavernosa. Graft was placed unstretched to prevent chordee (Figure 2). At the end of the procedure spongioplasty was done. Separated muscles are apposed and skin closed with 3-0 vicryl suture.



Figure 1: Buccal mucosa after being harvested.



Figure 2: Buccal mucosa placed as an onlay graft to the urethral plate. The patient remained bed rest for 72 hours and were discharged on the 5th to 7th postoperative days with catheter. All patients received broad-spectrum antibiotics and were kept on oral medications till removal of catheter. The catheter was left in situ for 3 weeks. After 3 weeks, a retrograde pericatheter urethrography was carried out and when no extravasation was found urethral catheter was removed. All patients were followed at 3 monthly intervals for one year and then six-monthly for 2 years and then annually. At each visit, patients were assessed for symptoms, urine culture and sensitivity, ultrasound to assess post void residual urine and uroflowmetry. The average follow up period of this study was 18

months.

Data were processed and analyzed using SPSS (Statistical Package for Social Sciences) software. The test statistics used to analyze the data were descriptive statistics, Chi-square test and student's t-Test. For all analytic tests, the level of significance was set at 0.05 and p value <0.05 was considered significant.

4. Results

A total of 40 patients (mean age 38 years, range 26 to 64 years) underwent dorsolateral onlay BMG urethroplasty between April 2017 to March 2020 in our hospital. Etiology of strictures, lichen sclerosus 19 cases (47.5%), Idiopathic 9 cases (22.5%), traumatic 6 cases (15%), catheter induced (reactive) inflammatory 4 cases (10%) and infective 2 cases (5%) (Table 1).

Most common presentation was poor urinary stream. Mean stricture length, as measured by preoperative RGU was 6.7 cm (range 2.3 cm to 13 cm). Site of the strictures were panurethral in 22 cases (55%), penile in 5 cases (12.5%), penobulbar in 7 cases (17.5%) and bulbar in 6 cases (15%).

Table 1: Etiology of strictures.

Etiology of stricture	Number of patients
Lichen sclerosus	19
Idiopathic	9
Traumatic	6
Catheter induced inflammatory	4
Infective	2
Total	40

Table 2: Site of stricture urethra

Site of stricture	Number of patients
Panurethral	22
Penile	5
Penobulbar	7
Bulbar	6
Total	40

Mean operative time was 180 minute (range 150 to 230 minute). Buccal mucosa was harvested from both the cheeks in 27 cases (67.5%) and from both cheeks and lower lip 13 cases (32.5%). Mean preoperative maximum flow rate (Qmax) was 5.87 ml/sec and mean postoperative maximum flow rate (Qmax) was 24.8 ml/sec. Pre operatively ultrasonography showed average post voided residual volume was 180. 4 ml which was significantly high. Post operatively average post voided residue was less than 25 ml 33 cases (82.5%). Five patients had narrowing at anastomotic site and presented with narrow urinary stream and prolonged voiding with abdominal straining. These patients were managed with Direct visual internal urethrotomy (DVIU). Two patients had developed acute urinary retention managed by suprapubic cystostomy and further urethroplasty procedure were required.

During postoperative period five patients (12.5%) developed mild wound infection, managed successfully with change in antibiotics as per culture sensitivity report and dressings. Two patients developed urethrocutaneous fistula and were managed conservatively. No patients developed diverticulum, sacculum formation, urinary

nary incontinence, erectile dysfunction and lower extremity injury.

Graft harvesting site complications: all patients experienced mild post-operative pain managed by good analgesia, facial swelling was reported five patients (12.5%). No patients encountered damage to Stensen's duct but transient lip paresthesia developed four patients (10%) and two patients (5%) reported restriction in mouth opening but resolved within one-month post operatively. No patients needed blood transfusion

5. Discussion

Urethral strictures are a frequent source of lower urinary tract disorders in adults. Urethral stricture disease has various etiologies. In the developed world post traumatic and post instrumentation are common causes of stricture. In developing country like India post inflammatory particularly balanitis xerotica obliterans is the most common cause of stricture and in our study also 47.5% of stricture were due to BXO and most of them were long segment and panurethral. Mean stricture length was 6.7 cm (range 2.3 cm to 13 cm). Most commonly the strictures were panurethral in our series which is comparable to other studies.

Substitution urethroplasty with BMG is most commonly performed technique particularly for long segment anterior segment stricture and become standard of care. In the dorsal onlay technique, the conventional circumferential dissection of the whole anterior urethra risks the vascularity of the urethra, which may be more important if the meatus is involved with the disease and the distal most urethra is extremely dissected, such is seen with concomitant balanitis xerotica obliterans. In dorsolateral onlay approach urethra is mobilized in one side, which maintain vascular integrity of urethra. It also preserves the one sided bulbar artery in addition to maintaining the native lateral vascularity in the meatus and distal urethra. Furthermore, by using this approach, we did not find post-operative chordee in any case. In this study, the urethra is not completely mobilized off the corpora; hence, graft sizing is more appropriate, preventing the chordee.

In this study total 40 patients underwent dorsolateral onlay BMG urethroplasty with overall success rate of 82.5%. Mean follow up was 18 months. Urethroplasty by Kulkarni et al shows study on 24 patients with mean follow up of 22 months shows 22(92%) had a successful outcome and 2(8%) were failure [10]. Studies by Barbagli or Kulkarni on dorsal onlay urethroplasty have shown success rate more than 90%.

In general, complications are rare after BMG urethroplasty. Post-operative complications can occur in two areas, the site of harvest and site of urethral stricture repair [11]. Potential buccal mucosal harvest site complications include haemorrhage, pain, facial swelling, damage to Stensen's duct, lip paresthesia, numbness and restriction in mouth opening. Facial swelling and restriction in mouth opening are common, but are self-limiting and will resolve within the first few months of surgery [12, 13] as reported

in our study. Wood et. al [14] reported that closure of the harvest site was associated with worse pain and suggested that this may be improved by not closing. Although Dublin et al. reported that patients did well with closure of the donor site, but 16% and 32% had long-term complaints of numbness and mouth tightness respectively. In our study we reported no difference of pain in two groups. Overall, the inner cheek donor site for BMG regardless of management appears to heal without complications as reported in our study [12, 13]. Perineal complications are similarly rare. Wound infection, haematomas, skin paresthesia, stricture and other local infections do sometimes occur. In a study by Fichtner J et al. they reported overall complication rate of 25%(8 of 32) [15]. In one more study overall complications rate was 5.4% with 5.7% restructure rate [12]. In our study we reported 19.8% overall complications rate which include restructure, wound complications and urethrocutaneous fistula.

6. Conclusion

Reconstructing the urethra continues to be challenging for the reconstructive surgeon. Buccal mucosal graft urethroplasty becomes standard care for long segment anterior urethral stricture. It is easy to harvest and to handle, is resilient to infections and accustomed to a wet environment. The preservation of one sided vascular supply to the urethra and its entire muscular and neurogenic support represents significantly towards perfecting the surgical technique of urethral reconstruction using a minimally approach like dorsolateral onlay BMG urethroplasty. In our study, we found that it is safe, technically feasible, easily adoptable, reproducible and finally successful procedure.

References

1. Lumen N, Hoebeke P, Willemsen P, de Troyer B, Pieters R, Oosterlinck W. Etiology of urethral stricture disease in the 21st century. *Journal of Urology*. 2009; 182(3): 983-7.
2. Berglund RK, Angermeier KW. Combined buccal mucosa graft and genital skin flap for reconstruction of extensive anterior urethral strictures. *Urology*. 2006; 68(4): 707-10.
3. Kumar MR, Himanshu A, Surdarshan O. Technique of anterior urethroplasty using the tunica albuginea of corpora cavernosa. *Asian Journal of Surgery*. 2008; 31(3): 134-9.
4. Barbagli G, Sansalone S, Djinovic R, Romano G, Lazzeri M. Current controversies in reconstructive surgery of anterior urethra: a clinical overview. *International Brazilian Journal of Urology*. 2012; 38(3): 307-16.
5. Barbagli G, Palminteri E, Guazzoni G, et al. Bulbar urethroplasty using buccal mucosa graft placed on the ventral, dorsal or lateral surface of the urethra: Are results affected by the surgical technique? *J Urol*. 2005; 174: 955-7.
6. Patterson JM, Chapple CR. Surgical techniques in substitution urethroplasty using buccal mucosa for the treatment of anterior urethral strictures. *EurUrol*. 2008; 53: 1162.
7. Singh BP, Pathak HR, Andankar MG. Dorsolateral onlay urethroplasty for anterior urethral strictures by a unilateral urethral mobilization approach. *Indian J Urol*. 2009; 25: 211-14.
8. Kulkarni BS, Barbagli G, Sansalone S, et al. One-sided anterior urethroplasty: a new dorsal onlay graft technique. *BJU Int*. 2009; 104: 1150-55.
9. Habib AKMK, Alam AKMK, Amanullah ATM, Rahman H, Hossain AKMS, Salam MA, Kibria SAMG. Dorsolateral onlay urethroplasty for long segment anterior urethral stricture: outcome of a new technique. 2011; 37(3): 78-82.
10. Kulkarni BS, Barbagli G, Sansalone S, et al. One sided anterior urethroplasty: a new dorsal onlay graft technique. *BJUI* 2009; 1-6.
11. Dublin N, Stewart LH. Oral complications after buccal mucosal graft harvest for urethroplasty. *BJU Int*. 2004; 94: 867-9.
12. Humby G. A one-stage operation for hypospadias repair. *Br J Surg*. 1941; 29: 84-92. Iselin CE, Webster GD. Dorsal onlay graft urethroplasty for repair of bulbar urethral stricture. *J Urol*. 1999; 161: 815-8.
13. Tolstunov L, Pogrel MA, McAninch JW. Intraoral morbidity following free buccal mucosal graft harvesting for urethroplasty. *Oral Surg Oral Med Oral Pathol Radiol Endod*. 1997; 84: 480-2.
14. Wood DN, Allen SE, Andrich DE, Greenwell TJ, Mundy AR. The morbidity of buccal mucosal graft harvest for urethroplasty and the effect of nonclosure of the graft harvest site on postoperative pain. *J Urol*. 2004; 172: 580-3.
15. Fichtner J, Filipas D, Fisch M, Hohenfellner R, Thuroff JW. Long-term outcome of ventral buccal mucosa onlay graft urethroplasty for urethral stricture repair. *Urol*. 2004; 64: 648-50.