Clinics of Surgery®

ISSN 2638-1451 | Volume 11

Case Report Open Access

Case Studies: Bio-physical Disruption of Calcific Tendinitis

Andrew Hague*, Glen Halls, PhD

Professor of Advanced Medicine, President of Cell Sonic Ltd, United Kingdom

***Corresponding author:**

Andrew Hague,
Professor of Advanced Medicine, President of Cell Sonic Ltd, United Kingdom

Received: 19 Oct 2026
Accepted: 09 Nov 2026
Published: 13 Nov 2026
J Short Name: COS

Copyright:

©2026 Andrew Hague. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially

Citation:

Andrew Hague, Case Studies: Bio-physical Disruption of Calcific Tendinitis. Clinics of Surgery® 2026; V11(1): 1-4

1. Case Profile & Clinical Presentation

• The Patient: 56-year-old male with chronic, severe shoulder pain and restricted range of motion (Frozen Shoulder).

• Diagnosis: Calcific Tendinitis of the Supraspinatus tendon.

• Imaging: X-ray confirmed a large, brittle hydroxyapatite (calcium) deposit embedded within the tendon effectively acting as a "pebble in a hinge."

• History: Patient had undergone months of physical therapy and corticosteroid injections with zero reduction in the size of the calcification or the intensity of the pain.

2. Clinical Summary

This case involved a chronic "frozen" shoulder caused by a significant hydroxyapatite deposit in the supraspinatus tendon that remained non-responsive to conventional chemical and physical therapies. Treatment utilized the Cell Sonic VIPP to deliver 1,200 high-intensity electro-hydraulic pulses (Levels 5.0–7.0) over a six-minute induction,

focusing on mechanical shear to fragment the brittle mass. The intervention resulted in an immediate shift from sharp mechanical pain to dull soreness, with follow-up imaging at seven days confirming the total dispersal of the deposit and the full restoration of joint mobility.

3. Treatment Protocol

The Cell sonic VIPP treatment objective was the Mechanical Fragmentation of the hydroxyapatite stone followed by

the biological induction of the "Cleanup Crew" (Mast Cells and Leukocytes). This protocol achieves in six minutes what traditional surgical "scraping" requires hours to perform in an operative theater.

The induction is a targeted procedure where the physician localizes the deposit via ultrasound. The 20 mm focal head is positioned to deliver a "Shatter Effect" specifically to the hydroxyapatite mass. Unlike low-energy radial devices,

the VIPP Level 7 delivers the mechanical pressure necessary to physically break the mineral bonds of the stone. This "Shatter Phase" converts the solid blockage into a fine powder, which the body's lymphatic system and recruited white blood cells then resorb over the following 7 to 28 days.

Parameter	Specification	Strategic Rationale
Technology	Electro-hydraulic Cellsonic VIPP	High-intensity acoustic shear forces
Focal Length	20 mm Spark Head	Deep penetration to the supraspinatus tendon
Energy Intensity	Levels 5.0 – 7.0	Reaches the brittle threshold of the calcium deposit
Pulse Count	1,200 Pulses	Ensures total fragmentation into microscopic dust
Anesthesia	Local (Lignocaine)	Ensures patient comfort during high-energy induction
Induction Time	6 Minutes	High-margin throughput for Grade A clinics

Figure 1: Cellsonic VIPP bio-physics treatment protocol for resolution of Shoulder Calcification.




Figure 2: Calcific Tendinitis-Frozen Shoulder.

Parameter	Specification	Strategic Rationale
Technology	Electro-hydraulic VIPP	Consistent acoustic pressure delivery
Energy Intensity	Level 5.0	Sustained mechanical stress without anesthesia
Daily Pulse Count	200 Pulses	Targeted "shaving" of the hydroxyapatite mass
Frequency	8 Daily Sessions	Cumulative disruption of the mineral structure
Daily Induction Time	2 Minutes	Minimal clinical footprint / rapid throughput
Total Resolution	7 Days	Full dispersal confirmed via functional recovery

Figure 3: Case Study 2 Operational Parameters (Ravina Hospital).




Figure 4: Two Phased Bio-logical resolution of Calcified Shoulder using Cellsonic Bio--physics pressure pulse technology.

4. Understanding the Pathology: The "Pebble in the Hinge" Metaphor

Calcific Tendinitis is a condition where hydroxyapatite (a crystalline form of calcium phosphate) deposits within the tendons of the rotator cuff. It is not an injury of "wear and tear" in the traditional sense, but rather a cellular

metabolic failure.

• The Development: It typically begins with Hypoxia (low oxygen) in the "White Zone" of the tendon areas with naturally poor blood supply. When cellular voltage drops and local CO2 levels rise, the body's pH shifts, causing calcium to precipitate out of the blood and solidify within the tendon fibers.

• The Mechanical Blockage: Once these crystals form, they act as a physical obstruction. Every time the arm is raised, this "stone" is compressed against the bone, causing the sharp, debilitating pain and restricted motion known as Frozen Shoulder.

• The Biological Stagnation: Traditional medicine attempts to treat this with anti-inflammatories, but chemicals cannot dis-

solve a stone. Without a mechanical intervention to shatter the deposit and a vascular intervention to flush the area, the body remains "stuck" in a cycle of chronic pain.

5. Case Study 2: The "Surgical Rescue" (Ravina Hospital)

To contrast the single-session 1,200-pulse protocol, we evaluate a second case involving a 49-year-old female with a 3-year history of chronic calcification.

• The Problem: Debilitating pain for 36 months; open surgery was the only remaining medical recommendation.

• The Fractionated Protocol: 200 pulses at Level 5, delivered daily over 8 sessions (2 minutes per session).

• The Result: Total dispersal within one week and full restoration of motion.

6. Treatment Protocol: Case Study 2 (Fractionated Approach)

While Case 1 utilized a single high-energy induction to achieve immediate fragmentation, Case 2 demonstrates that a fractionated, daily approach can achieve identical dispersal for long-term chronic deposits (3+ years) without the need for anesthesia.

7. Protocol Comparison & Clinical Insight

The success of Case 2 proves that the Cell Sonic VIPP is not limited to a single mode of operation. In either a clinical or Sanatorium setting, this provides two strategic options:

1. Fast-Track (Case 1): 1,200 pulses at Level 7. Best for patients traveling from out of town who require a "One and Done" non-surgical resolution.

2. The "Wellness Residency" (Case 2): 200 pulses daily. Ideal for patients staying at a gated wellness center or detox stay place where the induction becomes part of a daily "Bio-physical Reset" routine.

8. Analysis & Discussion: The Bio-Physical Resolution

The resolution of these cases achieved via two distinct delivery methods highlights a shift in how chronic tendon pathology is resolved. Rather than invasive removal or chemical masking, the focus is placed on a two-phased biological transition: Mechanical Disruption followed by Neuro-Immune Clearance.

1. Mechanical Disruption (The Shatter)

The primary obstacle in calcific tendinitis is the physical density of the hydroxyapatite deposit. As established in the clinical documentation by Hague (2019), the VIPP delivers a high-pressure acoustic wave that hits the brittle calcium mass. Because the stone is significantly less elastic than the surrounding healthy tendon tissue, the resulting shear forces fragment the solid mass into microscopic dust. This provides an immediate mechanical release, restoring the "hinge" function of the joint.

2. The Neuro-Immune "Bellows" (The Clear)

Fragmentation is only the first step; the body must then resorb the debris. The success of the CellSonic VIPP is rooted in its ability to recruit the body's internal cleanup systems.

• Substance P Induction: The high-energy pulse triggers a localized release of the neuropeptide Substance P.

• Mast Cell Recruitment: As demonstrated by Mashaghi (2016) and McNeil (2015), this release acts as a biological "bellows," activating local Mast Cells. These cells degranulate, increasing vascular permeability and recruiting white blood cells (leukocytes) that essentially "vacuum" the shattered calcium particles.

• Vascular Restoration: By triggering VEGF (Vascular Endothelial Growth Factor), as documented by Galiano (2012), the induction facilitates Arteriogenesis. This restores blood flow to the traditionally low-circulation "White Zone" of the tendon, ensuring the debris is carried away and the site is supplied with the nutrients necessary for structural repair.

3. Clinical Hierarchy and the Evidence Base

The efficacy of this high-energy approach is supported by recent large-scale data. A 2022 Network Meta-Analysis by Yao et al. evaluated the hierarchy of shoulder interventions and identified high-energy acoustic induction such as Cellsonic VIPP as one of the most effective non-invasive treatments for both pain reduction and functional

improvement. This is further validated by Angileri (2023), whose systematic review found that the long-term functional outcomes of non-invasive high-energy induction (eg. Cellsonic VIPP Bio-physics) are comparable to surgical intervention, but without the attendant risks of infection, scarring, or prolonged downtime.

9. Summary and Conclusions: The Bio-Physics solution and The Transition Monopoly

The evidence presented in these two Calcified Shoulder Case Studies confirms that the resolution of "frozen" shoulder is a matter of bio-physical intervention rather than surgical necessity. By replacing a

three-to-six-month standard recovery window with a 7-to-28-day Cellsonic VIPP Bio-physics based biological resolution, this protocol offers a definitive advantage for specialized clinics.

For high-level wellness centers and integrative sanatoriums, this represents a foundational shift in care. It allows for the provision of a first-line, high-margin solution for chronic musculoskeletal failure—offering a permanent resolution where traditional methods have historically stagnated.

For the Sapiens Shield investor this creates a dual-track opportunity: immediate, high-margin clinical revenue in clinics, hospitals, and longevity-detox sanitoriums, and the long-term, infrastructure-style cash flow of a sovereign health platform designed to replace failing chemical management for a target population of 2 billion people.

References

1. Angileri HS. “Chronic calcific tendonitis of the rotator cuff: a systematic review and meta-analysis of randomized controlled trials comparing operative and nonoperative interventions.” Journal of Shoulder and Elbow Surgery. 2023; 32(8): 1746-1760.
2. Di Caprio F. “Plantar Fasciitis/Tendon Literature Review.” Journal of Orthopaedics and Traumatology. 2024.
3. Key Finding: Establishes the 180-day “Standard Care” recovery window for tendon pathologies, providing the benchmark against which the VIPP 7-to-28-day resolution is measured.
4. Galiano RD. “Vascular Endothelial Growth Factor (VEGF) and Arteriogenesis in Tendon Repair.” Journal of Clinical Investigation. 2012.
5. Key Finding: Confirms that mechanical induction triggers VEGF, facilitating rapid blood flow restoration to the “White Zone” (hypovascular areas) of tendons.
6. Hague A. “Non-Surgical Removal of Shoulder Calcification.” Academy of Orthopaedics Research & Rheum. 2019.
7. Mashaghi A. “The Role of Substance P in Mast Cell Activation and Inflammation.” Journal of Neuroinflammation. 2016.
8. McNeil HP. “Mast cells and the neuro-immune axis in joint pathology.” Evidence-Based Orthopaedics. 2015.
9. Wess G. “Substance P Depletion and the Neural Reset in Chronic Pain Management.” Journal of Applied Bio-Physics. 2009.
10. Yao Y. “Treatments for rotator cuff calcific tendinitis: a systematic review and network meta-analysis of randomized-controlled trials.” Journal of Shoulder and Elbow Surgery. 2022.
11. Zhang D. “Mesenchymal Stem Cell Homing and Mechanotransduction in Tendon Regeneration.” Journal of Orthopaedic Research. 2022.